



MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Minimal Model Program and Adjunction Theory

Marco Andreatta

Dipartimento di Matematica di Trento, Italia

Moscow, October 2015



Quasi polarized pairs

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Let X be a projective variety of dimension n with good singularities, for example **terminal and \mathbb{Q} -factorial singularities**.

Let L be a **Cartier divisor** (a line bundle) which is **ample**, or simply **nef and big**.

The pair (X, L) is called a **polarized pair**, or a **quasi polarized pair**.

Example: $X \subset \mathbb{P}^N$ be a projective variety and $L := \mathcal{O}(1)|_X$, or better its (partial) desingularization and the pull back of L .



Classical problems

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castelnuovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$,
MMP

$K_X + \Delta^{(n-2)}$,
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Problem Given a general element $D \in |L|$
(assume that X is not a cone over D).

Which properties of D lift to X ;
do these properties determine X ?

Enriques–Castelnuovo studied the case in which X is a surface and D is a curve of low genus, or of minimal degree, ...

Fano studied the case in which X is a 3-fold and D is a $K3$ surface.

Mori in his first paper proved that if D is a c.i. in a weighted projective space the same is for X .

Sommese proved that abelian and bi-elliptic surfaces **cannot be ample sections**, unless X is a cone.



Weighted Blow-up

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_Y + \Delta^{(n-1)}$.
MMP

$K_Y + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Let $\sigma = (a_1, \dots, a_n) \in \mathbb{N}^{+n}$ such that $\gcd(a_1, \dots, a_n) = 1$.

The weighted projective space with weight (a_1, \dots, a_n) is defined as:

$$\mathbb{P}(a_1, \dots, a_n) := (\mathbb{C}^n - \{0\})/\mathbb{C}^*,$$

where $\xi \in \mathbb{C}^*$ acts by $\xi(x_1, \dots, x_n) = (\xi^{a_1}x_1, \dots, \xi^{a_n}x_n)$.

A cyclic quotient singularity,

$$\mathbb{C}^n/\mathbb{Z}_m(a_1, \dots, a_n) := X,$$

is an affine variety definite as the quotient of \mathbb{C}^n by the action $\epsilon(x_1, \dots, x_n) = (\epsilon^{a_1}x_1, \dots, \epsilon^{a_n}x_n)$, where $\epsilon \in \mathbb{Z}_m$.



Weighted Blow-up

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Consider the rational map

$$\varphi : \mathbb{C}^n / \mathbb{Z}_m(a_1, \dots, a_n) \rightarrow \mathbb{P}(a_1, \dots, a_n)$$

given by $(x_1, \dots, x_n) \mapsto (x_1 : \dots : x_n)$.

Definition

The weighted blow-up of $X = \mathbb{C}^n / \mathbb{Z}_m(a_1, \dots, a_n)$ with weight σ , \bar{X} , is defined as the closure in $X \times \mathbb{P}(a_1, \dots, a_k)$ of the graph of φ , together with the morphism $\pi_\sigma : \bar{X} \rightarrow X$ given by the projection on the first factor.



Weighted Blow-up

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$,
MMP

$K_X + \Delta^{(n-2)}$,
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

For any $d \in \mathbb{N}$ we define the σ -weighted ideal of degree d as

$$I_{\sigma,d} = \{g \in \mathbb{C}[x_1, \dots, x_n] : \sigma\text{-wt}(g) \geq d\} = (x_1^{s_1} \cdots x_n^{s_n} : \sum_{j=1}^n s_j a_j \geq d).$$

Proposition

The weighted blow-up of $X = \mathbb{C}^n / \mathbb{Z}_m(a_1, \dots, a_n)$ with weight σ , $\pi : \bar{X} \rightarrow X$, is given by

$$\bar{X} = \text{Proj}_X \left(\bigoplus_{d \geq 0} I_{\sigma,d} \right) \rightarrow X.$$



Lifting Weighted Blow-Ups

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_Y + \Delta^{(n-1)}$.
MMP

$K_Y + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Theorem

Let $f : X \rightarrow Z$ be an elementary local projective contraction with \mathbb{Q} -factorial singularities, which contracts a divisor E to a point $p \in Z$. Assume $\dim X \geq 4$.

Let $Y \subset X$ be a f -ample Cartier divisor and assume that $f' =: f_Y : Y \rightarrow f(Y) := W$ is a $\sigma' = (a_1, \dots, a_{n-1})$ -blow-up.

Then $f : X \rightarrow Z$ is a $\sigma = (a_1, \dots, a_{n-1}, a_n)$ -blow-up, $\pi_\sigma : X \rightarrow Z$, $a_n = a$, where a is the positive integer such that $Y|_{E \cap Y} = \mathcal{O}_{\mathbb{P}}(a)$.

Remark

*The **Cox Rings** of weighted blow-ups determines completely the blow-up (it is Toric). The following could be a good question: does the Cox Ring of an ample section determine the one of the variety?*



Proof

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

By assumption Z has an isolated \mathbb{Q} -factorial singularity and $W \subset Z$ is a Weil divisor which has a cyclic quotient singularity, i.e.

$$W = \mathbb{C}^{n-1} / \mathbb{Z}_m(a_1, \dots, a_{n-1}).$$

We first prove that $Z = \mathbb{C}^n / \mathbb{Z}_m(a_1, \dots, a_{n-1}, a_n)$.

Then, let b a positive integer such that $-bE \sim Y$ (and therefore Cartier).

By Grothendieck theory $X = \text{Proj}_{\mathcal{O}_Z}(\oplus_{d \geq 0} f_*(\mathcal{O}_X(-dbE)))$, therefore we want to prove that

$$f_*(\mathcal{O}_X(-dbE)) = I_{\sigma, db} = (x_1^{s_1} \cdots x_n^{s_n} \mid \sum_{j=1}^n a_j s_j \geq db).$$



Proof

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$,
MMP

$K_X + \Delta^{(n-2)}$,
MMP

$\binom{n-3}{n-2} < \tau \leq$

Consider the exact sequence on X

$$0 \rightarrow \mathcal{O}_X(-Y - dbE) \rightarrow \mathcal{O}_X(-dbE) \rightarrow \mathcal{O}_Y(-dbE) \rightarrow 0$$

Pushing it down via φ and using the relative Vanishing theorems we have

$$0 \rightarrow f_* \mathcal{O}_X(-(d-1)bE) \xrightarrow{\cdot x_n} f_* \mathcal{O}_X(-dbE) \rightarrow f_* \mathcal{O}_Y(-dbE) \rightarrow 0.$$

The proposition follows by **induction on n**

$$(f_*(\mathcal{O}_Y(-dbE) = (x_1^{s_1} \cdots x_{n-1}^{s_{n-1}} \mid \sum_{j=1}^{n-1} a_j s_j \geq db)))$$

and on d

$$(f_*(\mathcal{O}_X(-(d-1)bE) = (x_1^{s_1} \cdots x_n^{s_n} \mid \sum_{j=1}^n a_j s_j \geq (d-1)b))$$



Adjunction Theory

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Adjunction Theory wants to classify quasi polarized pairs via the study of the nefness of the adjoint bundles

$$K_X + rL,$$

with r natural (or rational) positive number.

Assume that there exist r sections of $|L|$ which intersect in a $n - r$ variety D , with terminal singularities (if $r = n - 1$, we ask for a smooth curve).

To get nefness of $K_X + rL$ implies, by adjunction $(K_X + rL)|_D = K_D$, to get a **minimal model for D** .



Minimal Model Program- BCHM

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

A log pair (X, Δ) , i.e a normal variety X and an effective \mathbb{R} divisor Δ , is **Kawamata log terminal (klt)** if:

$K_X + \Delta$ is \mathbb{R} -Cartier and for a (any) log resolution $g : Y \rightarrow X$ we have $g^*(K_X + \Delta) = K_Y + \sum b_i \Gamma_i$ with $b_i < 1$, for all i .

By BCHM on a klt log pair (X, Δ) , with Δ big, we can run a

$K_X + \Delta$ - Minimal Model Program with scaling:

$$(X_0, \Delta_0) = (X, \Delta) \rightarrow (X_1, \Delta_1) \rightarrow \cdots \rightarrow (X_s, \Delta_s)$$



Minimal Model Program- BCHM

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

$(X_0, \Delta_0) = (X, \Delta) \rightarrow (X_1, \Delta_1) \rightarrow \cdots \rightarrow (X_s, \Delta_s)$
such that:

- 1) (X_i, Δ_i) is a klt log pair, for $i = 0, \dots, s$;
- 2) $\varphi_i : X_i \rightarrow X_{i+1}$ is a birational map which is either a **divisorial contraction** or a **flip** associated with an **extremal ray** $R_i = \mathbb{R}^+[C_i]$ such that $(K_{X_i} + \Delta_i) \cdot C_i < 0$
(notation: $R_i \in \overline{NE(X_i)}_{(K_{X_i} + \Delta_i) < 0} \subset \overline{NE(X_i)}_{K_{X_i} < 0}$)
- 3) either $K_{X_s} + \Delta_s$ is nef (i.e. (X_s, Δ_s) is a **log Minimal Model**), or $X_s \rightarrow Z$ is a **Mori fiber space relatively to $K_{X_s} + \Delta_s$**
(depending on the pseudoeffectivity of $K_X + \Delta$).



MMP for a q.p. pair

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ^r -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\binom{n-3}{n-2} < \tau \leq$

Let (X, L) be a quasi-polarized variety and let $r \in \mathbb{Q}^+$.

Lemma (zip L into a boundary). Since L is nef and big there exists an effective \mathbb{Q} -divisor Δ^r on X such that

$$rL \sim_{\mathbb{Q}} \Delta^r \quad \text{and} \quad (X, \Delta^r) \text{ is Kawamata log terminal.}$$

Run a **$K_X + \Delta^r$ -MMP** and get a birational klt pair (X_s, Δ_s^r) which is

- either a Minimal Model ($K_{X_s} + \Delta_s$ is nef)
- or $X_s \rightarrow Z$ is a Mori fiber space relatively to $K_{X_s} + \Delta_s$.

Remarks/Problems

- (X_s, Δ_s^r) is **not necessarily an (r) q.p. pair**, i.e. we do not have a priori a nef and big Cartier divisor L_s such that $rL_s \sim_{\mathbb{Q}} \Delta_s^r$.
- Beyond the **existence** of the MMP, it would be nice to have a **"description"** of each steps and eventually of the Mori fiber spaces.



Extremal rays

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$,
MMP

$K_X + \Delta^{(n-2)}$,
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

For the above program we study the (Fano-Mori) contractions:

$$\varphi : X \rightarrow Y$$

associated to a rays $R = \mathbb{R}^+[C] \in \overline{NE(X)}_{(K_X+rL)<0} \subset \overline{NE(X)}_{K_X<0}$.

That is φ is a projective map between normal variety, with connected fibers, X has terminal \mathbb{Q} -factorial singularities and an irreducible curve $C \subset X$ is mapped to a point by φ iff $[C] \in R$.

- φ can be of fiber type ($\dim X > \dim Y$), φ is called a Mori fiber space
- or birational, φ can then be divisorial or small

Let F be a non trivial fiber of φ ; we possibly restrict to an affine neighborhood of the image of F (local set up).

Then $(K_X + \tau L) \sim_{\varphi} \mathcal{O}_X$ for a rational $\tau > r$.



Apollonio method

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Let $X' \in |L|$ a generic divisor with "good singularities". We have:

- $\varphi|_{X'} := \varphi' : X' \rightarrow Y'$ is a contraction with connected fibre,
around $F' := F \cap X'$;

it is the Fano-Mori contraction associate to $R' \in \overline{NE(X')}_{(K_{X'} + (r-1)L') < 0}$.

- Any section of L on X' extends to a section of L on X .



Base point free technique

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Theorem [Fano, Fujita, Kawamata, Kollar, Shokurov, ...,
..., A-Wisniewski, Mella, A-Tasin]

- $\dim F > (r - 1)$; if φ is birational then $\dim F > r$.
- If $\dim F < r + 1$, or $\dim F \leq r + 1$ if φ is birational, then L is very ample (relatively to φ).
- If $\dim F < r + 2$ then there exists $X' \in |L|$ with "good" singularities (i.e. as in X). The same is true if $\dim F = r + 2$, except for two cases in which $n = 3$ and φ is of fiber type.



Castelnuovo-Kawakita

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castelnuovo-Kawakita contractions

$K_X + \Delta^{(n-1)}$ -MMP

$K_X + \Delta^{(n-2)}$ -MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Theorem

Let $\varphi : X \rightarrow Y$ be a birational contraction in a $K_X + \Delta^{n-2}$ -MMP) (i.e. it is associated with an extremal ray on a q.p. pair such that $R = \mathbb{R}^+[C] \in \overline{NE(X)}_{(K_X+(n-2)L)<0} \subset \overline{NE(X)}_{K_X<0}$ and $L \cdot C > 0$).

Then $\varphi : X \rightarrow Y$ is the weighted blow-up of a smooth point in Y of weights $(1, 1, b, \dots, b)$, where b is a natural positive number.

$L' = \varphi_*(L)$ is a Cartier divisor on Y .

Definition

We call such φ a Castelnuovo-Kawakita contraction.



Proof

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

We have $\dim F > (n-2)$; thus $\dim F = (n-1)$ and φ is a contraction of a divisor to a point.

By the base point free theorem, we can assume L is very ample.

Thus we get the existence of sections in $|L|$ with terminal singularities.

Inductively, slicing with $(n-2)$ general sections of $|L|$, we can reduce to the case of a Fano Mori contraction on a surface, $f' : S \rightarrow W$.

Surfaces with terminal singularities are smooth. Apply now

Castelnuovo's Theorem to have that W is smooth and f' is a $(1,1)$ -blow-up.

Apply the "lifting of weighted blow-up".



$$r = (n - 1)$$

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Consider a $K_X + \Delta^r$ -MMP with $r = (n - 1)$ (or $\geq (n - 1)$) and let $R_i = \mathbb{R}^+[C_i]$ be a birational ray in the sequence.

Inductively construct a nef and big Cartier divisor L_i on X_i such that $rL_i \sim_{\mathbb{Q}} \Delta_i^{(n-1)}$:

1) $L_i C_i = 0$, otherwise, by the above Theorem, we have the contradiction $(n - 1) \geq \dim F > r \geq (n - 1)$.

2) Let $\varphi_i : X_i \rightarrow Y$ be the contraction associated with R_i . We have a Cartier divisor L'_{i+1} such that $\varphi_i^*(L'_{i+1}) = L_i$.

If φ is birational $(X_{i+1}, L_{i+1}) := (Y, L'_{i+1})$,

if φ is small $(X_{i+1}, L_{i+1}) := (X_i^+, \varphi^+(L'_{i+1}))$, where $\varphi^+ : X_i^+ \rightarrow Y$ is the flip.



the zero reduction

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Proposition. Given a q.p. pair (X, L) it is possible to run a MMP which contracts **all** extremal rays on which L is zero and obtain a q.p. pair (X', L') which is birational equivalent to (X, L) and such that:

- either $K_{X'} + (n-1)L'$ is nef
- or (X', L') is a Mori space relative to $K_{X'} + (n-1)L'$ and L' is a (relatively) very ample Cartier divisor.

Definition. (X', L') is called a **zero reduction** of (X, L) .

By very **classical results** in the second case the q.p. pair (X', L') is in a obvious finite list of examples: $(\mathbb{P}^n, \mathcal{O}(1))$, $(Q, \mathcal{O}(1))$, scrolls, del Pezzo.



Applications

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Let (X, L) be a quasi-polarized variety and $g(X, L)$ be its sectional genus: $2g(X, L) - 2 = (K_X + (n-1)L) \cdot L \cdots L$.

(if L is spanned it is the genus of a curve intersection of $n-1$ general elements in $|L|$).

- Classification of pairs $g(X, L) \leq 0$ ($K_X + (n-1)L$ is not nef (therefore not pseudoeffective) and the zero reduction of (X, L) is among the above pairs

- Classification of pairs with $g(X, L) = 1$.

- Classification of pairs of minimal degree
(i.e. $L^n = h^0(X, L) - n$).



First Reduction

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castelnuovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Proposition-Part 1. Let (X, L) be a q.p. pair. There exists a q.p. pair (X'', L'') which is a $(K_X + \Delta^{n-2})$ -MM and which can be obtained with the following procedure:

- Take a zero reduction (X', L') .
- Contract, step by step, all Castelnuovo-Kawakita type extremal rays, such that $L'_{|E} = -bE_{|E}$; $\varphi' : X' \rightarrow X''$.
- Let $L'' := \varphi'_* L'$.

Definition The pair (X'', L'') is called a **First Reduction** of the pair (X, L) .



First Reduction

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Proposition-Part 2. Let (X, L) be a q.p. pair and let (X'', L'') be its First Reduction. Then

- either $K_{X''} + (n-2)L''$ is nef
- or $X'' \rightarrow Z$ is a Mori fiber space relatively to $K_{X''} + (n-2)L''$ and L'' is (relatively) very ample with one exception (del Pezzo manifold). In all cases there exists a divisor in $|L''|$ with good singularities.

Remark. The classification of the pairs in the second part, thank to the existence of a good section, is classical and reduces to the theory of algebraic surfaces. (Quadric fibration, del Pezzo manifolds,)



Applications

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Theorem. Let $Y \subset \mathbb{P}^N$ be a non degenerate projective variety of dimension $n \geq 3$ of degree d and let $\tilde{L} := \mathcal{O}(1)|_Y$. Assume that $d < 2\text{codim}_{\mathbb{P}^N}(X) + 2$.

Then on a desingularization (X, L) the divisor $K_X + (n-2)L$ is not pseudoeffective.

Therefore $(Y, \mathcal{O}(1))$ is equivalent, via birational equivalence and first-reduction, to a q.p. pair (X'', L'') in the above Remark.



$$(n-3) < \tau \leq (n-2)$$

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$,
MMP

$K_X + \Delta^{(n-2)}$,
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

Let $\varphi : X \rightarrow Y$ be a birational contraction associated with an extremal ray $R = \mathbb{R}^+[C]$ on a q.p. pair, such that $L \cdot C > 0$ and $(n-3) < \tau \leq (n-2)$.

(These are the birational maps in a $K_X + \Delta^{n-3}$ -MMP).

We have the following possibilities for φ .

- 1 φ contracts a divisor to a curve (it is a special case of the next theorem),
- 2 φ contracts a divisor to a point,
- 3 φ is a small contraction with exc. locus of dimension $(n-2)$.

In all cases we can apply the "base point free technique", find therefore a section $X' \in |L|$ and, by induction if possible, reduce to the case $n = 3$ where we have a complete classification.

Then we like to "lift" this classification and the examples to higher dimension.



Divisorial contractions with small fibers

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$(n-3) < \tau \leq (n-2)$

Theorem. Let $f : X \rightarrow Z$ be a divisorial contraction associated to an extremal ray R in $\overline{NE}(X)_{(K_X + rL) < 0}$, where $r \in \mathbb{N}^+$, such that $L \cdot C > 0$. Let E be the exceptional locus of f and set $C := f(E) \subset Z$.

Assume that all fibres have dimension less or equal to $r + 1$.

- 1 Then there is a closed subset $S \subset Z$ of codimension at least 3 such that $Z' = Z \setminus S$ and $C' = C \setminus S$ are smooth, $\text{codim}_{Z'} C' = r + 2$ and $f' : X' = X \setminus f^{-1}(S) \rightarrow Z'$ is a weighted blow-up along C' with weight $\sigma = (1, 1, b, \dots, b, 0, \dots, 0)$, where the number of b 's is r .
- 2 Let \mathcal{I}' be a σ -weighted ideal sheaf of degree b for $Z' \subset X'$ and let $i : Z' \rightarrow Z$ be the inclusion; let also $\mathcal{I} := i_*(\mathcal{I}')$ and $\mathcal{I}^{(m)}$ be the m -th symbolic power of \mathcal{I} . Then $X = \text{Proj } \bigoplus_{m \geq 0} \mathcal{I}^{(m)}$.



Kawakita Contractions

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_Y + \Delta^{(n-1)}$.
MMP

$K_Y + \Delta^{(n-2)}$.
MMP

$\frac{(n-3)}{(n-2)} < \tau \leq$

The second case was treated for $n = 3$ in a series of paper by Kawakita. By the Theorem on lifting weighted blow-up we can extend Kawakita classification and show that φ is a weighted blow-up of a (possible singular) point.



Small Contractions

MMP on q. p.v.

Marco Andreatta

Polarized variety

Weighted Blow-up

Δ' -MMP

Extremal rays

Castenovo-
Kawakita
contractions

$K_X + \Delta^{(n-1)}$.
MMP

$K_X + \Delta^{(n-2)}$.
MMP

$\binom{n-3}{n-2} < \tau \leq$

The third case was treated for $n = 3$ by S. Mori and by S. Mori and J. Kollar.

At the moment we can extend to higher dimension the Francia's flip, i.e. the case where X has only points of index one and two.