A differential criterium for regularity of quaternionic functions

Alessandro Perotti¹

*Dipartimento di Matematica, Università degli Studi di Trento, Via Sommarive 14, 38050 Povo-Trento, Italy
Fax: +39 0461881624, perotti@science.unitn.it*

Abstract

Let $\Omega \subseteq \mathbb{C}^2$. We prove that there exist differential operators T and N , with complex coefficients, such that a function $f : \overline{\Omega} \rightarrow \mathbb{H}$ of class C^1 is regular if and only if $(N - jT)f = 0$ on $\partial\Omega$ (j a basic quaternion) and f is harmonic on Ω . At the same time we generalize a result of Kytmanov and Aizenberg. We show that a complex harmonic function h on Ω ($\partial\Omega$ connected) is holomorphic if and only if $\bar{\partial}_n h = \overline{aL(h)}$ on $\partial\Omega$, where $\bar{\partial}_n$ is the normal part of $\bar{\partial}$, L is a tangential Cauchy-Riemann operator and $a \in \mathbb{C}$.

1. Introduction

Let $\Omega = \{z \in \mathbb{C}^2 : \rho(z) < 0\}$ be a bounded domain in $\mathbb{C}^2$ with boundary of class $C^m, m \geq 1$. We assume $\rho \in C^m$ on $\mathbb{C}^2$ and $d\rho \neq 0$ on $\partial\Omega$. Let ν denote the outer unit normal to $\partial\Omega$ and $\tau = i\nu$. For every $F \in C^1(\overline{\Omega})$, we set $\bar{\partial}_n F = \frac{1}{2} \left(\frac{\partial F}{\partial \nu} + i \frac{\partial F}{\partial \tau} \right)$ (see [1] §§3.3 and 14.2). In a neighbourhood of $\partial\Omega$ we have the decomposition of $\bar{\partial}F$ in the tangential and the normal parts: $\bar{\partial}F = \bar{\partial}_b F + \bar{\partial}_n F \frac{\bar{\partial}\rho}{|\bar{\partial}\rho|}$.

The normal part of $\bar{\partial}F$ on $\partial\Omega$ can also be expressed by means of the Hodge $*$ -operator and the Lebesgue surface measure $d\sigma$, as $\bar{\partial}_n F d\sigma = *\bar{\partial}F|_{\partial\Omega}$. We denote by L the tangential Cauchy-Riemann operator

$$L = \frac{1}{|\bar{\partial}\rho|} \left(\frac{\partial \rho}{\partial \bar{z}_2} \frac{\partial}{\partial \bar{z}_1} - \frac{\partial \rho}{\partial \bar{z}_1} \frac{\partial}{\partial \bar{z}_2} \right).$$

Let $\mathbb{H}$ be the algebra of quaternions $q = x_0 + ix_1 + jx_2 + kx_3$, where x_0, x_1, x_2, x_3 are real numbers and i, j, k denote the basic quaternions. We identify the space $\mathbb{C}^2$ with the set $\mathbb{H}$ by means of the mapping that associates to $(z_1, z_2) = (x_0 + ix_1, x_2 + ix_3)$ the quaternion $q = z_1 + z_2 j$. The commutation rule is

Email address: perotti@science.unitn.it (Alessandro Perotti).

URL: www.science.unitn.it/~perotti (Alessandro Perotti).

¹ Partially supported by MIUR (Project “Proprietà geometriche delle varietà reali e complesse”) and GNSAGA of INdAM.

then $aj = j\bar{a}$ for every $a \in \mathbb{C}$. We refer to [9] for the basic facts of quaternionic analysis. We will denote by D the left Cauchy-Riemann-Fueter operator

$$D = \frac{\partial}{\partial x_0} + i \frac{\partial}{\partial x_1} + j \frac{\partial}{\partial x_2} + k \frac{\partial}{\partial x_3}.$$

A quaternionic C^1 function f is *(left-)regular* on a domain Ω if $Df = 0$ on Ω . We will use also another class of regular functions, which are in the kernel of the following differential operator defined by the structural vector $\psi = \{1, i, j, -k\}$:

$$D' = \frac{\partial}{\partial x_0} + i \frac{\partial}{\partial x_1} + j \frac{\partial}{\partial x_2} - k \frac{\partial}{\partial x_3} = 2 \left(\frac{\partial}{\partial \bar{z}_1} + j \frac{\partial}{\partial \bar{z}_2} \right).$$

A quaternionic C^1 function $f = f^1 + f^2 j$, is called *(left-) ψ -regular* on a domain Ω if $D'f = 0$ on Ω . This condition is equivalent to the following system of complex differential equations for the complex components (f^1, f^2) of f :

$$\frac{\partial f^1}{\partial \bar{z}_1} = \frac{\partial \bar{f}^2}{\partial \bar{z}_2}, \quad \frac{\partial f^1}{\partial \bar{z}_2} = -\frac{\partial \bar{f}^2}{\partial \bar{z}_1} \quad (1)$$

or to the equation $*\bar{\partial}f^1 = -\frac{1}{2}\bar{\partial}(\bar{f}^2 d\bar{z}_1 \wedge d\bar{z}_2)$. Note that the $(1, 2)$ -form $*\bar{\partial}f^1$ is $\bar{\partial}$ -closed on Ω when f^1 is harmonic. Then, if Ω is a pseudoconvex domain in $\mathbb{C}^2$, every complex harmonic function f^1 on Ω is a complex component of a ψ -regular function $f = f^1 + f^2 j$. (cf. [5] and [7] for this result and its converse). *Remark 1 Any holomorphic mapping (f^1, f^2) on Ω defines a ψ -regular function $f = f^1 + f^2 j$. Moreover, the complex components of a ψ -regular function are either both holomorphic or both not-holomorphic.*

We refer, for instance, to [8], [4] or [6] for the properties of structural vectors and ψ -regular functions. Regular and ψ -regular functions are real analytic on Ω , and they are harmonic with respect to the Laplace operator in $\mathbb{R}^4$.

Remark 2 Let γ be the transformation of $\mathbb{C}^2$ defined by $\gamma(z_1, z_2) = (z_1, \bar{z}_2)$. Then a C^1 function f is regular on the domain Ω if, and only if, $f \circ \gamma$ is ψ -regular on $\gamma^{-1}(\Omega)$.

2. A differential criterium for regularity

Theorem 2.1 *Let Ω be a bounded domain in $\mathbb{H}$, with smooth boundary of class C^1 . Let $f = f^1 + f^2 j : \bar{\Omega} \rightarrow \mathbb{H}$ be a function of class C^1 . Then, f is (left-) ψ -regular on Ω if, and only if, f is an harmonic function on Ω and*

$$(\bar{\partial}_n - jL)f = 0 \quad \text{on } \partial\Omega. \quad (2)$$

Proof If f is a ψ -regular function on Ω , then from equations (1) we get that it satisfies the equation $(\bar{\partial}_n - jL)f = 0$ on $\partial\Omega$, since $\bar{\partial}_n f^1 = -\bar{L}(f^2)$, $\bar{\partial}_n f^2 = \bar{L}(f^1)$ on $\partial\Omega$.

Conversely, assume that $\bar{\partial}_n f = jL(f)$ on $\partial\Omega$. For every $z \in \Omega$, it follows from the Bochner-Martinelli integral representation for complex harmonic functions on Ω (see for example [1]§1.1), that $f(z) = f^1(z) + f^2(z)j$ is equal to

$$\int_{\partial\Omega} U(\zeta, z) f^1(\zeta) + 2 \int_{\partial\Omega} g(\zeta, z) \bar{\partial}_n f^1(\zeta) d\sigma + \int_{\partial\Omega} U(\zeta, z) f^2(\zeta) j + 2 \int_{\partial\Omega} g(\zeta, z) \bar{\partial}_n f^2(\zeta) j d\sigma$$

where $U(\zeta, z)$ is the two-dimensional Bochner-Martinelli form and $g(\zeta, z) = \frac{1}{4\pi^2} |\zeta - z|^{-2}$ is the fundamental solution of the complex laplacian on $\mathbb{C}^2$. Then we obtain

$$f(z) = \int_{\partial\Omega} U(\zeta, z) f(\zeta) + 2 \int_{\partial\Omega} g(\zeta, z) \bar{\partial}_n f(\zeta) d\sigma = \int_{\partial\Omega} U(\zeta, z) f(\zeta) + 2 \int_{\partial\Omega} g(\zeta, z) jL(f(\zeta)) d\sigma.$$

In [10] (see also [4]) it was shown that the two-dimensional Bochner-Martinelli form can be considered as a first complex component of the Cauchy-Fueter kernel $G'(p - q)$ associated to ψ -regular functions. Let $q = z_1 + z_2 j$, $p = \zeta_1 + \zeta_2 j$, $\sigma(q) = dx[0] - idx[1] + jdx[2] + kdx[3]$, where $dx[k]$ denotes the product of dx_0, dx_1, dx_2, dx_3 with dx_k deleted. Then $G'(p - q)\sigma(p) = U(\zeta, z) + \omega(\zeta, z)j$, where $\omega(\zeta, z)$ is the complex (1, 2)-form $\omega(\zeta, z) = -\frac{1}{4\pi^2} |\zeta - z|^{-4} ((\bar{\zeta}_1 - \bar{z}_1)d\zeta_1 + (\bar{\zeta}_2 - \bar{z}_2)d\zeta_2) \wedge d\bar{\zeta}$. From Proposition 6.3 in [4], we get that

$$f(z) = \int_{\partial\Omega} U(\zeta, z) f(\zeta) + \int_{\partial\Omega} \omega(\zeta, z) j f(\zeta) = \int_{\partial\Omega} G'(p - q)\sigma(p) f(p).$$

The Cauchy-Fueter integral formula gives the ψ -regularity of f .

Let N and T be the differential operators, defined in a neighbourhood of $\partial\Omega$,

$$N = \frac{\partial\rho}{\partial z_1} \frac{\partial}{\partial \bar{z}_1} + \frac{\partial\rho}{\partial \bar{z}_2} \frac{\partial}{\partial z_2}, \quad T = \frac{\partial\rho}{\partial z_2} \frac{\partial}{\partial \bar{z}_1} - \frac{\partial\rho}{\partial \bar{z}_1} \frac{\partial}{\partial z_2}.$$

T is a tangential (not Cauchy-Riemann) operator to $\partial\Omega$, while N is non-tangential, such that $N(\rho) = |\bar{\partial}\rho|^2$, $\text{Re}(N) = |\bar{\partial}\rho| \text{Re}(\bar{\partial}_n)$. The remark 2 shows that Theorem 2.1 gives also a boundary condition for regularity of a harmonic function on Ω .

Corollary 2.2 *Let Ω be a C^1 -bounded domain in $\mathbb{H}$. Let $f = f^1 + f^2 j : \bar{\Omega} \rightarrow \mathbb{H}$ be a function of class C^1 . Then, f is (left-)regular on Ω if, and only if, f is harmonic on Ω and*

$$(N - jT)f = 0 \quad \text{on } \partial\Omega.$$

3. A differential criterium for holomorphicity in $\mathbb{C}^2$

Theorem 3.1 *Let Ω be a bounded domain in $\mathbb{C}^2$, with connected boundary of class C^1 . Let $a \in \mathbb{C}$. If $h \in C^1(\bar{\Omega})$ is complex harmonic on Ω and satisfies the differential condition $\bar{\partial}_n h = \overline{aL(h)}$ on $\partial\Omega$, then h is holomorphic on Ω .*

Proof We use the same notation of the proof of theorem 2.1. Let F^+ and F^- be the ψ -regular functions defined respectively on Ω and on $\mathbb{C}^2 \setminus \bar{\Omega}$ by the Cauchy-Fueter integral of $f = ah + hj$:

$$F^\pm(z) = \int_{\partial\Omega} U(\zeta, z) f(\zeta) + \int_{\partial\Omega} \omega(\zeta, z) j f(\zeta).$$

From the equalities $U(\zeta, z) = -2*\partial_\zeta g(\zeta, z)$, $\omega(\zeta, z) = -\partial_\zeta(g(\zeta, z)d\bar{\zeta})$, we get that

$$\overline{F^-(z)} = -2 \int_{\partial\Omega} (\overline{ah(\zeta)} - j\overline{h(\zeta)}) * \bar{\partial}_\zeta g(\zeta, z) + \int_{\partial\Omega} (jah(\zeta) + h(\zeta)) \bar{\partial}_\zeta(g(\zeta, z)d\zeta)$$

for every $z \notin \bar{\Omega}$. By application of the Stokes' Theorem, of the complex Green formula and of the equality $\bar{\partial}h \wedge d\zeta|_{\partial\Omega} = 2L(h)d\sigma$ on $\partial\Omega$ we see that the equation $\bar{\partial}_n h = \overline{aL(h)}$ on $\partial\Omega$ implies that

$$\int_{\partial\Omega} \overline{h(\zeta)} * \bar{\partial}_\zeta g(\zeta, z) = -\frac{a}{2} \int_{\partial\Omega} h(\zeta) \bar{\partial}_\zeta(g(\zeta, z)d\zeta).$$

Therefore, F^- is a complex-valued, ψ -regular function on $\mathbb{C}^2 \setminus \overline{\Omega}$. The equations (1) then imply that F^- is a holomorphic function. Since $\partial\Omega$ is connected, F^- can be holomorphically continued to the whole space. Let $\tilde{F}^-$ be such extension. Then $F = F^+ - \tilde{F}^-|_{\Omega}$ is a ψ -regular function on Ω , continuous on $\overline{\Omega}$, such that $F|_{\partial\Omega} = f|_{\partial\Omega}$. This means that the pair (ah, h) is equal on Ω to the two complex components (f^1, f^2) of a ψ -regular function. From equations (1) we get that h satisfies

$$a \frac{\partial h}{\partial \bar{z}_1} = \overline{\frac{\partial h}{\partial \bar{z}_2}}, \quad \bar{a} \frac{\partial h}{\partial \bar{z}_2} = -\frac{\partial h}{\partial \bar{z}_1}$$

from which it follows $\bar{\partial}h = 0$.

Remark 3 The result stated in theorem 3.1 generalizes the following result of Kytmanov and Aizenberg [3], which holds in $\mathbb{C}^n$, $n > 1$: if h is a complex harmonic function on Ω , of class C^1 on $\overline{\Omega}$, such that $\partial_n h = 0$ on $\partial\Omega$, then h is holomorphic. See also [2] for a similar generalization.

References

- [1] A.M. Kytmanov, The Bochner-Martinelli integral and its applications, Birkhäuser Verlag, Basel, 1995
- [2] A.M. Kytmanov, Some differential criteria for the holomorphy of functions in $\mathbb{C}^n$, in: Some problems of multidimensional complex analysis, Akad. Nauk SSSR Sibirsk. Otdel., Inst. Fiz., Krasnoyarsk, 1980, pp. 51–64, 263–264 (Russian)
- [3] A.M. Kytmanov and L.A. Aizenberg, The holomorphy of continuous functions that are representable by the Bochner-Martinelli integral, Izv. Akad. Nauk Armyan. SSR 13 (1978) 158–169 (Russian)
- [4] I.M. Mitelman and M.V. Shapiro, Differentiation of the Martinelli-Bochner integrals and the notion of hyperderivability, Math. Nachr. 172 (1995) 211–238
- [5] M. Naser, Hyperholomorphe Funktionen, Sibirsk Mat. Zh. 12 (1971) 1327–1340 (Russian); English transl. in Siberian Math. J. 12 (1971) 959–968 (1972)
- [6] K. Nōno, α -hyperholomorphic function theory, Bull. Fukuoka Univ. Ed. III 35 (1985) 11–17
- [7] K. Nōno, Characterization of domains of holomorphy by the existence of hyper-conjugate harmonic functions, Rev. Roumaine Math. Pures Appl. 31 (1986) no.2 159–161
- [8] M.V. Shapiro and N.L. Vasilevski, Quaternionic ψ -hyperholomorphic functions, singular integral operators and boundary value problems. I. ψ -hyperholomorphic function theory, Complex Variables Theory Appl. 27 (1995) no.1 17–46
- [9] Sudbery, Quaternionic analysis, Mat. Proc. Camb. Phil. Soc. 85 (1979) 199–225
- [10] N.L. Vasilevski and M.V. Shapiro, Some questions of hypercomplex analysis, in: Complex analysis and applications '87 (Varna, 1987), Publ. House Bulgar. Acad. Sci., Sofia, 1989, pp. 523–531